

BMJ home	Current issue	Past issues	Classified ads	Career focus	Feedback
	Collections	About this site	About the BMJ	BMA	Medline

Statistics at Square One

Answers to exercises

1.1 Median 0.71, range 0.10 to 1.24, first quartile 0.535, third quartile 0.84 $\mu\text{mol}/24\text{ hr}$

2.1 Mean = 2.41, SD = 1.27.

2.2 Mean = 0.697 $\mu\text{mol}/24\text{ hr}$, SD = 0.0214 $\mu\text{mol}/24\text{ hr}$, range . 0.215 to 1.179 $\mu\text{mol}/1$

2.3 Points 0.10 and 1.24. 2/40 or 5%.

3.1 SE(mean) = 0.074 $\mu\text{mol}/24\text{ hr}$

3.2 A uniform or flat distribution. Population mean 4.5, population SD 2.87.

3.3 The distribution will be approximately Normal, mean 4.5 and SD $287/\sqrt{5} = 1.28$.

4.1 The reference range is 12.26 - 57.74, and so the observed value of 52 is included in it.

4.2 95% CI 32.73 to 37.27.

5.1 0.42 g/dl, $z = 3.08$ $0.001 < P < 0.01$, difference = 1.3 g/dl, 95% CI 0.48 to 2.12 g/dl.

5.2 0.23 g/dl, $P < 0.0001$.

6.1 SE (percentage) = 2.1%, SE (difference) = 3.7%, difference = 3.4%. 95% CI -3.9 to 10.7%, $z = 0.94$, $P = 0.35$.

6.2 Yes, the traditional remedy, $z = 2.2$, $P = 0.028$.

7.1 37.5 to 40.5 KA units.

7.2 $t = 2.652$, d.f. = 17, $0.01 < P < 0.02$.

7.3 0.56g/dl, $t = 1.243$, d.f.=20, $0.1 < P < 0.05$, 95% CI -0.38 to 1.50g/dl.

7.4 15 days, $t = 1.758$, d.f. = 9, $0.1 < P < 0.05$, 95% CI -4.30 to 34.30 days.

8.1 Standard $\chi^2 = 23.295$, d.f. = 4, $P > 0.5$. Trend $\chi^2 = 2.25$, d.f. = 1, $P = 0.13$.

8.2 $\chi^2 = 3.916$, d.f. = 1, $0.02 < P < 0.05$, difference in rates 9%, 95% CI 0.3 to 17.9%.

8.3 $\chi^2 = 0.931$, d.f. = 1, $0.1 < P < 0.5$, difference in rates 15%, 95% CI -7.7 to 38%.

8.4 $\chi^2 = 8.949$, d.f. = 3, $0.02 < P < 0.05$. Yes, practice C; if this is omitted the remaining practices give $\chi^2 = 0.241$, d.f. = 2, $P > 0.5$. (Both χ^2 tests by quick method.)

9.1 Sickness rate in first department 28%, in second department 8%, difference 20% (approximate 95% CI = -6 to 45%, $P = 0.24$ (Fisher's Exact test mid P)). P is calculated from $2 \times (0.5 \times 0.173 + 0.031)$.

10.1 Smaller total = -30. No.

10.2 Mann-Whitney statistic = 74. The group on the new remedy. No.

11.1 $r = -0.848$.

11.2 $r_s = -0.867$.

11.3 $y = 36.1 - 2.34x$. This means that, on average, for every 1 mile increase in mean distance the attendance rate drops by 2.34%. This can be safely accepted only within the area measured here.

11.4 SE = 0.39, 95% CI = $-2.34 - 2.145 \times 0.39$ to $-2.34 + 2.145 \times 0.39 = -3.1$ to -1.5% .

12.1 $O_A = 6$, $t_A = 8.06$, $O_B = 8$, $E_B = 5.94$. Log rank $\chi^2 = 1.24$, d.f. = 1, $0.1 < P < 0.5$.

12.2 Risk = 0.55, 95% CI 0.19 to 1.60.

13.1 Matched case control study.

13.2 Cohort study.

13.3 Cross sectional study.

13.4 Randomised controlled trial.

13.5 Quasi experimental design.